



**KNOX
MOUNTAIN
DENTISTRY**



#7 – 740 Clement Ave.
Kelowna BC V1Y 0H7



(778) 721-0011



hello@knoxmountaindentistry.com



@knoxmountaindentistry

Dental Extraction Post-Operative Instructions

The initial healing period usually takes one to two weeks. The following instructions will aid in avoiding postoperative discomfort and complications.

Immediately Following Procedure

- Encourage clotting by keeping steady biting pressure on the gauze placed at the bleeding site by your doctor. Pressure helps reduce bleeding and permits formation of a clot in the tooth socket. Fresh gauze has been provided to you, replace the gauze when it becomes saturated. Gently remove the gauze after the local anesthesia has worn off and normal feeling has returned (one to two hours is common).
- Some oozing of blood may persist after 24 hours. If necessary, apply gauze or a moist tea bag to the area and bite firmly for one hour. If bleeding does not slow, call the Clinic or the dentist who performed the extraction.
- Avoid activities that could apply a suction action to the blood clot such as sucking through a straw, spitting, swishing and sneezing with a closed mouth. All tobacco products should be stopped for 5 days following extractions to prevent healing problems. If tobacco use cannot be avoided, place a rolled piece of gauze over the extraction site and take gentle, short drags of the cigarette to minimize the possibility of clot dislodgement.
- Reduce your activity level for several hours minimum. Be careful not to bite or burn your mouth or lips. Avoid eating and unnecessary talking for a few hours.
- Do not rinse your mouth or brush your teeth for 24 hours. These activities may hinder formation of a blood clot, which is necessary for proper healing.

Control Pain and Swelling

- Swelling is normal after teeth have been extracted and should not cause alarm (during certain procedures, particularly wisdom tooth extraction, your doctor may administer an anti-swelling medication at the time of surgery). Swelling is at its greatest during the first 48 hours and will normally decline after this period. If swelling increases after 48 hours, call us.
- Immediately following the procedure, begin taking medication as directed by your doctor to minimize discomfort. Take it before the local anesthetic wears off for optimal pain control. Continue to take medication while pain persists.
- For mild discomfort take Ibuprofen (400-600mg) or Tylenol (500-1000mg) with soft food every four to six hours. For severe pain take the pharmaceutical drug as instructed-be sure to read the directions carefully and do not exceed the maximum daily dose of any medication. **In most cases, effective management of moderate pain is achieved with 1000mg Tylenol AND 600mg ibuprofen taken together every 6 hours for the first several days. Do not continue this regimen for longer than 5 days or as instructed by your doctor.**

- Use ice packs on surgical area (side of face) for 24 - 48 hours: apply ice 10 minutes on - 10 minutes off. Bags of frozen peas work well. Be sure to place a soft cloth (such as a wash cloth) between your face and the cold pack to avoid skin irritation. A prone position, with the head turned to one side, will allow you to lay the ice pack on to the sore muscle. Flip your body over and apply the ice pack to the other side of your face if needed for reduction of swelling and pain.

24 Hours After Surgery

- After bleeding has stopped, cautiously begin your daily mouth cleaning routine, but avoid disturbing the surgical site so as not to loosen or remove the blood clot.
- Brush your teeth twice per day and floss once per day. Brush the tongue with a wet toothbrush to keep bacteria growth down.

48 Hours After Surgery

- Rinsing is important to prevent infection and promote healing because it removes food particles and debris from the extraction site. Keeping your mouth very clean will help avoid infections, complications and delayed healing. Be gentle when rinsing, no vigorous swishing.
- Use a warm salt-water rinse (1/2 teaspoon salt per 8 oz cup of water) to clean the surgical area. Repeat after every meal or snack for seven days minimum or as long as you need to keep the area free of debris.
- Call our office immediately if you develop a rash, vomit, faint, or develop a temperature over 102 degrees.

Nutrition and Hydration

- Drink plenty of fluids. (Do not use a straw!)
- Diet may consist of soft foods, which can be easily chewed and swallowed. No seeds, nuts, rice, popcorn, hot beverages, alcohol, etc. Have your meals at the usual time. Eat what you wish, but be careful not to disturb the blood clot. Add solid foods to your diet as soon as they are comfortable to chew.

Sutures

- Occasionally stitches, or sutures, are placed in the gums, especially if more than one tooth is removed. Although you may feel them with your tongue, it is best to leave them alone. Gut sutures will dissolve in 2-5 days. If silk or synthetic suture material is used, the dentist will remove them in 5-7 days. The space left by the tooth will feel a bit strange to you at first. Eventually, new bone and gum tissue will grow into the gap left by the extraction.

If you have any questions or concerns, do not hesitate to contact the office at (778) 721-0011 or Dr. Sanders' cell (250) 307-7757.

